

泰国蛛毛苣苔属一新种*

许兆然

(中山大学生物系, 广州 510275)

A NEW SPECIES OF *PARABOEIA* RIDLEY FROM THAILAND*

Xu Zhao-ran

(Biology, Department, Zhongshan University, Guangzhou 510275)

Key words *Paraboeia*; *P. burttii* Z. R. Xu

关键词 蛛毛苣苔属; 布氏蛛毛苣苔

布氏蛛毛苣苔 新种

Paraboeia burttii Z. R. Xu, sp. nov. Fig. 1

P. capitatae Ridley proxima, sed foliis supra pubescentibus, bracteis valde majoribus, sepalis obovatis et majoribus differt.

Herba perennis rosulata, brevi-caulis vel acaulis; caules ad 15 cm longi 7 mm in diam. interdum internodiis < 1.5 cm longis. Folia opposita petiolata; petiolus 4—10 cm longus lanatus; lamina ovata, c. $7-14 \times 3-8$ cm, apice obtusa, basi cuneata vel plus minusve cordata, margine sparse-serrulata vel serrata, supra pubescens, subtus implicito-lanata nervis supra laevigatis subtus prominulis, lateralibus 6—10-jugis, venulis retem formantibus Inflorescentia axillaris, cymosa, floribunda, aspectu capitato vel umbellato, indumento implicito-lanato tecta praeter corollam, pedunculo 13—18 cm longo, bracteis plurimis subrotundis c. 1. 2 cm longis, pedicellis 1—1.5 cm longis. Flos c. 16 mm in diam. Calyx 5-sectus ad basim, Lanatus extra, glaber intra, sepalis obovatis c. 6×4.5 mm. Corolla glabra, c. 12 mm longa, 16 mm diam., tubo c. 4 mm longo, limbo 2-labiato, labio gupero 2-lobato, lobis ovatis c. 6×10 mm, labio infero 3-lobato, lobo medio obovato c. 4×6 mm, lobis lateralibus late ovatis c. 8×9 mm. Stamina 2 glabra, antheris reniformibus c. 2.5×4 mm apice

* The author is grateful to Royal Botanic Garden, Edinburgh and Sibbold Trust for their funding and facility supports to this study, and would like to thank Mr. B. L. Burtt and Mr. Xie QingJian for the former's academic guidance and the latter's assistance in making the line drawing in this work.

1993-01-03 收稿。

cohaerentibus gynoeceum versus ad medium introrsum apertis, dorsifixis, filamentis brevibus ad basim corollae adnatis. Gynoeceum glabrum, c. 10 mm longum, ovario ovoideo c. 3×2 mm, stylo c. 7 mm longo, stigmate stylo leviter latiore c. 1×1.3 mm. Capsula linearis, 3—4 cm longa, 1.5 mm in diam. glabra, haud convoluta, horizontaliter effusa, seminibus ovoideis c. 0.4×0.3 mm.

Thailand: Nakhawan Srithamarat, Khiriwong, Khao Khi Naw., herb common by stream in evergreen forest, flowers bluish, 5 Oct 1957 (fr), Bunnak 710 (holotype, E);




图1 布氏蛛毛苣苔

Fig. 1 *Paraboea burttii* Z. R. Xu new sp.

Wam Tai, 20 km N of Trang, 11 Oct 1970 (fr), C. Charoenphol *et al.* 3640 (E); cultivated in Aberdeen, UK, 1930? (fl), A. F. G. Kerr 205 (L, K.); Bangkok, 2 Sep 1930 (fl), A. F. G. Kerr 19745 (K); Pattalung, Tamote Distr., Tamote Falls National Park., 10 Sep 1986 (fl), J. F. Maxwell 86-651 (PSU); Nakornsitamarat, Lansagah, 14 Sep 1985 (fl), J. F. Maxwell 877 (A); Distr. Lansagah Gahrome Falls, Khao Luang National Park, 13 Dec 1985 (fr), J. F. Maxwell 1101 (A, PSU), 14 Sep 1985 (fl), J. F. Maxwell 85877 (PSU); Rao Chem, Tung Sang, 20 Jul 1929 (ster), Rabil 103 (K), (vernacular name; Dardhoi) 20 Jul 1929 (fl), Rabil 106 (K); Khao Luang National Park, 23 Aug 1980 (fl), P. Sirirugsa 328 (PSU); Nakawn Sritamarat, Kiriwong, 26 Jul 1951 (fr), T. Smitinand 710 (BKF, J); Khao Rawn Nai Hawn, 27 Nov 1951 (fl), P. Sunarnakoses 193 (BKF); hill above Ronpibun, 16 Jul 1940 (fl), F. K. Ward 27460 (K).

本种看起来和 *P. capitata* Ridley 相近, 但后者的苞片少而且小, 花蕾期没有包裹花序; 另外, 萼片小而且卵圆形, 易于区别。模式标本产泰国。

苦苣苔科蛛毛苣苔属 *Paraboea* Ridley 共84种(其中中国产13种), 分布在中国南部直至婆罗洲和印度; 绝大部分种类产于石灰岩石上, 多数种类蒴果螺旋形扭转, 并以叶背具有一层致密的毛被为特征。Burtt 于1984年发表该属的初步订正, 把叶背具有 *arachnoid indumentum* (蛛丝状毛) 作为该属特征。潘开玉编写《中国植物志》的同属植物时, 采用 Burtt 的处理, 并根据 Burtt 提出的 *arachnoid indumentum* 新拟中文属名为“蛛毛苣苔属”。根据 Burtt 和我近年的研究, 这种毛被并不是严格的蛛丝状毛, 而是比较密实的山羊绒状毛被 (*matted cashmere-like indumentum*)。这层毛被覆盖在叶子下表面, 把气孔口全部荫蔽, 以减少水分从气孔蒸腾而损失, 是植物一种典型的抗旱特征。野外工作时不难看到锈色蛛毛苣苔 *P. rufescens* 常常生长在毫无荫蔽的裸露石灰岩石上。我们研究这个属的专著将于近期在爱丁堡皇家植物园出版。